



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

readily obtained. In numerical problems the task of solving the transcendental equations (7) or (10), and (7a) presents no practical difficulty.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

172. Proposed by H. C. FEEMSTER, York, Neb.

Show that $\frac{(nr)!}{n!(r!)^n}$ is an integer.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

By induction we get the following:

Suppose $r(n-1)!$ is divisible by $(r!)^{n-1}(n-1)!$. Now

$$\begin{aligned} \frac{(rn)!}{n!(r!)^n} &= \frac{r(n-1)!}{(r!)^{n-1}(n-1)!} \times \frac{(nr)!}{r(n-1)!} \times \frac{(n-1)!}{n!r!} \\ &= \frac{nr(nr-1)(nr-2)\dots \text{to } r \text{ factors}}{nr(r-1)!} \\ &= \frac{(nr-1)(nr-2)\dots \text{to } (r-1) \text{ factors}}{(r-1)!} = \text{an integer.} \end{aligned}$$

Now $\frac{r!}{r!1!} = \text{an integer}$, and hence $\frac{(2r)!}{(r!)^2 2!} = \text{an integer}$; $\frac{(3r)!}{(r!)^3 3!} = \text{an integer}$, and so on.

173. Proposed by V. M. SPUNAR, M. and E. E., East Pittsburg, Pa.

Find integral values satisfying the equation,

$$a_1^2 + a_2^2 + a_3^2 + \dots + a_n^2 = d^4.$$

I. Solution by FRANK LOXLEY GRIFFIN, Ph. D., Williams College.

One set of solutions may be obtained by putting $a_1 = df_1$, $a_2 = df_2$, etc., which reduces the problem to that of finding a sequence of n integers, the sum of whose squares is a perfect square. Or, geometrically, we seek $n-1$ right triangles whose sides are all integers, and the hypotenuse of each being one leg of the next.

I. This is readily accomplished for $n=2$ by recalling that, if $(p^2 - q^2)$ and $2pq$ are legs, where p and q are integers, the hypotenuse is also an integer, $(p^2 + q^2)$. Let f_1 be *any odd integer* (>1) and take $p - q = 1$ and $p + q = f_1$, so that $p = \frac{1}{2}(f_1 + 1)$, $q = \frac{1}{2}(f_1 - 1)$, both integers. Thus the sides

of the right triangle are f_1 , $\frac{1}{2}(f_1^2-1)$, and $\frac{1}{2}(f_1^2+1)$. Or, calling the second f_2 and the third d , we have $d^2=f_1^2+f_2^2$.

II. Let this value of d for $n=2$ be denoted by d_2 , and observe that d_2 is odd. For f_1 =say, $2m+1$ (being odd), whence $d_2=2m^2+2m+1$, odd. Thus d_2 may be treated precisely as was f_1 , giving $f_3=\frac{1}{2}(d_2^2-1)$, $d_3=\frac{1}{2}(d_2^2+1)$, or $d_3^2=d_2^2+f_3^2=f_1^2+f_2^2+f_3^2$.

Since again d_3 is odd, the process may clearly be continued indefinitely. The successive values of f and d may be found from: $2f_{n+1}=d_n^2-1$, $2d_{n+1}=d_n^2+1$. The following table gives the values up to $n=6$ if we begin with f_1+3 :

f_n	3	4	12	84	3612	6526884
d_n		5	13	85	3613	6526885

To obtain the proposed α 's for any value of n , merely multiply each f (up to f_n) by d_n . Thus, for $n=3$,

$$39^2+52^2+156^2=13^4.$$

[We might also start with any *even* integer (>2), and either (A) form a sequence all terms of which are even multiples of the terms obtaining by beginning with any odd factor of f_1 , or (B) let $f_1=2pq$ and take for p, q any factors whose product is $\frac{1}{2}f_1$. Thus for $f_1=14$:

$$\begin{array}{llll} p=7, & q=1, & f_2=48, & d_2=50, \\ p_2=25, & q_2=1, & f_3=624, & d_3=626, \text{ etc.} \end{array}$$

II. Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Dr. Martin's solution can be extended to suit this requirement.

Let $x_1^2+x_2^2+x_3^2+\dots+x_n^2=c^2$.

Then $c^2-n^2=(c-n)(c+n)=x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2$.

Let $c-n=b$, then $c+n=(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2)/b$.

$\therefore x_n=\frac{1}{2}(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2-b^2)/b$.

$c=\frac{1}{2}(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2+b^2)/b$.

Multiplying through by c^2 after substituting, we get

$$\begin{aligned} (cx_1)^2+(cx_2)^2+(cx_3)^2+\dots+[(c/2b)(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2-b^2)]^2 \\ =[(1/2b)(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2+b^2)]^4. \end{aligned}$$

$$\therefore (4b^2cx_1)^2+(4b^2cx_2)^2+(4b^2cx_3)^2+\dots[2bc(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2-b^2)]^2=(x_1^2+x_2^2+x_3^2+\dots+x_{n-1}^2+b^2)^4.$$

Let $n=7$, $b=3$, $x_1, x_2, x_3, \dots=1, 2, 3, \dots, 6$. Then $c=\frac{5}{3}$.

$$\therefore (600)^2+(1200)^2+(1800)^2+(2400)^2+(3000)^2+(3600)^2+(8200)^2=(100)^4, \text{ or } 6^2+12^2+18^2+24^2+30^2+36^2+82^2=10^4.$$

Also solved by the Proposer.